

Histopathology of the pancreas in fulminant type 1 diabetes after 23-year follow-up: a case report.

Sato T, Miki T, Murakami N, Hirohashi Y, Kouzu H, Furuhashi M, Tanno M, Yuda S, Saitoh S, Hasegawa T, Miura T.

Pathol Int. 2012;62:827-9.

Synergistic prognostic values of cardiac sympathetic innervation with left ventricular hypertrophy and left atrial size in heart failure patients without reduced left ventricular ejection fraction: a cohort study.

Doi T, Nakata T, Hashimoto A, Yuda S, Wakabayashi T, Kouzu H, Kaneko N, Hase M, Tsuchihashi K, Miura T.

BMJ Open. 2012;2.

Cytoprotective regulation of the mitochondrial permeability transition pore is impaired in type 2 diabetic Goto-Kakizaki rat hearts.

Itoh T, Kouzu H, Miki T, Tanno M, Kuno A, Sato T, Sunaga D, Murase H, Miura T.

J Mol Cell Cardiol. 2012;53:870-9.

Trials, tribulations and speculation! Report from the 7th Biennial Hatter Cardiovascular Institute Workshop.

Bell R, Beeuwkes R, Bøtker HE, Davidson S, Downey J, Garcia-Dorado D, Hausenloy DJ, Heusch G, Ibanez B, Kitakaze M, Lecour S, Mentzer R, Miura T, Opie L, Ovize M, Ruiz-Meana M, Schulz R, Shannon R, Walker M, Vinten-Johansen J, Yellon D.

Basic Res Cardiol. 2012;107:300.

Effects of parental hypertension on longitudinal trends in blood pressure and plasma metabolic profile: mixed-effects model analysis.

Mitsumata K, Saitoh S, Ohnishi H, Akasaka H, Miura T.

Hypertension. 2012;60:1124-30.

Role of ER stress in ventricular contractile dysfunction in type 2 diabetes.

Takada A, Miki T, Kuno A, Kouzu H, Sunaga D, Itoh T, Tanno M, Yano T, Sato T, Ishikawa S, Miura T.

PLoS One. 2012;7:e39893.

Elevation of fatty acid-binding protein 4 is predisposed by family history of hypertension and contributes to blood pressure elevation.

Ota H, Furuhashi M, Ishimura S, Koyama M, Okazaki Y, Mita T, Fuseya T, Yamashita T, Tanaka M, Yoshida H, Shimamoto K, Miura T.

Am J Hypertens. 2012;25:1124-30.

Effects of diabetes on myocardial infarct size and cardioprotection by preconditioning and postconditioning.

Miki T, Itoh T, Sunaga D, Miura T.

Cardiovasc Diabetol. 2012;11:67.

Emerging beneficial roles of sirtuins in heart failure.

Tanno M, Kuno A, Horio Y, Miura T.

Basic Res Cardiol. 2012;107:273.

Role of connexin-43 in protective PI3K-Akt-GSK-3 β signaling in cardiomyocytes.

Ishikawa S, Kuno A, Tanno M, Miki T, Kouzu H, Itoh T, Sato T, Sunaga D, Murase H, Miura T.

Am J Physiol Heart Circ Physiol. 2012;302:H2536-44.

Cardiac mortality assessment improved by evaluation of cardiac sympathetic nerve activity in combination with hemoglobin and kidney function in chronic heart failure patients.

Doi T, Nakata T, Hashimoto A, Yuda S, Wakabayashi T, Kouzu H, Kaneko N, Hase M, Tsuchihashi K, Miura T.

J Nucl Med. 2012;53:731-40.

Diabetic cardiomyopathy: pathophysiology and clinical features.

Miki T, Yuda S, Kouzu H, Miura T.

Heart Fail Rev. 2012 Mar.

A single nucleotide polymorphism of the adenosine deaminase, RNA-specific gene is associated with the serum triglyceride level, abdominal circumference, and serum adiponectin concentration.

Oguro R, Kamide K, Katsuya T, Akasaka H, Sugimoto K, Congrains A, Arai Y, Hirose N, Saitoh S, Ohishi M, Miura T, Rakugi H.

Exp Gerontol. 2012;47:183-7.

Fusion of bone marrow-derived cells with renal tubules contributes to renal dysfunction in diabetic nephropathy.

Yamashita T, Fujimiya M, Nagaishi K, Ataka K, Tanaka M, Yoshida H, Tsuchihashi K, Shimamoto K, Miura T.

FASEB J. 2012;26:1559-68.

Angiotensin II receptor activation in youth triggers persistent insulin resistance and hypertension--a legacy effect?

Togashi N, Maeda T, Yoshida H, Koyama M, Tanaka M, Furuhashi M, Shimamoto K, Miura T.

Hypertens Res. 2012;35:334-40.